



1. **STANDARDIZATION**
 2. **STANDARDIZATION**

ROBBER COMMERCIAL	PUNTO BLANCO		
REPRESENTANTE LEGAL	CATTIA SERRA-PATINO PERAZZO		
CONTADOR	DAVID RODRIGUEZ JUAN CARLOS		
CLASE CONTINENTE	STACA	DELIADO LLIVAS CONTABILIDAD	SI
CALIFICACIÓN APTESADAL	SI	SI/NO	SI

HPLC column:		Flow rate:	1.0 mL/min
HPLC solvent:	CH ₂ Cl ₂ /MeOH	Dye:	Acridine orange
HPLC column temperature:		Excitation wavelength:	360 nm

ACTIVATING OR DEACTIVATING A USER

[illegible]

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----



Journal of Management Education 34(1)

- **SECTOR PRIVADO:** INSTITUCIONES SOCIALES, EMPRESAS DEL SECTOR Y ASOCIACIONES
- **SECTOR PÚBLICO:** MINISTERIO DE TRANSPORTES Y OBRAS PÚBLICAS
- **SECTOR RELACIONADO:** COMERCIO
- **SECTOR TURISMO:** COMISIÓN DE TURISMO
- **SECTOR DE LA EDUCACIÓN:** LA LECTURA, SOCIEDADES
- **SECTOR DE LA SALUD:** EN LA PRÁCTICA
- **SECTOR DE LA CULTURA:** EN LA PRÁCTICA

IN THE DISTRICT COURT OF THE UNITED STATES FOR THE DISTRICT OF COLUMBIA	IN RE: THE ESTATE OF	ADMINISTRATOR
JOHN W. WILSON	JOHN W. WILSON	JOHN W. WILSON



Abstract: *Chlamydia trachomatis* is the most common bacterial sexually transmitted infection in the United States. The purpose of this study was to determine the prevalence of *C. trachomatis* in a community-based sample of men and women in the United States. A cross-sectional study was conducted in 1999 in a community-based sample of men and women in the United States. The prevalence of *C. trachomatis* was 1.1% (95% CI 0.4-2.8%) in men and 1.1% (95% CI 0.4-2.8%) in women. The prevalence of *C. trachomatis* was significantly higher in men than in women (p = 0.001). The prevalence of *C. trachomatis* was significantly higher in men than in women (p = 0.001).

Revised: 2004-01-17 10:40:45 PM



STAGNANT TROOP
FURNITURE, FURNISHINGS, FURNITURE CO.



[illegible]

Copyright © 2006 John Wiley & Sons, Ltd.

Page 28/40