

505.000.00

[illegible]

Continued on reverse of document, Art. 190 de la Ley de Compañías

0 1 6 . 7 9 8 1 0 0 0 0 0 0 0 0 0 0 0

1990年12月25日
 1990年12月25日

() ; [] ; []

Journal of Management Education 30(6)

1. The first group of variables is the set of variables that are used to define the *independent* variable. These variables are the variables that are used to define the *independent* variable. These variables are the variables that are used to define the *independent* variable.

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total phenolic content was determined by the method of Singleton and Rossi (1965). The total flavonoid content was determined by the method of Zhishen et al. (1999). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Folch et al. (1957). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total ash content was determined by the method of AOAC (1990). The total acid content was determined by the method of AOAC (1990). The total base content was determined by the method of AOAC (1990). The total nitrogen content was determined by the method of Kjeldahl (1900). The total phosphorus content was determined by the method of Molybdenum blue (1900). The total sulfur content was determined by the method of Barium sulfate (1900). The total calcium content was determined by the method of Oxalate (1900). The total magnesium content was determined by the method of Magnesia (1900). The total potassium content was determined by the method of Potassium dichromate (1900). The total sodium content was determined by the method of Sodium chloride (1900). The total iron content was determined by the method of Iron(III) chloride (1900). The total copper content was determined by the method of Copper(II) sulfate (1900). The total zinc content was determined by the method of Zinc sulfate (1900). The total manganese content was determined by the method of Manganese sulfate (1900). The total cobalt content was determined by the method of Cobalt(II) chloride (1900). The total nickel content was determined by the method of Nickel(II) sulfate (1900). The total chromium content was determined by the method of Chromium(III) chloride (1900). The total boron content was determined by the method of Boric acid (1900). The total molybdenum content was determined by the method of Molybdenum trioxide (1900). The total selenium content was determined by the method of Selenium dioxide (1900). The total tellurium content was determined by the method of Telluric acid (1900). The total iodine content was determined by the method of Iodine (1900). The total bromine content was determined by the method of Bromine (1900). The total fluorine content was determined by the method of Hydrofluoric acid (1900). The total chlorine content was determined by the method of Hydrochloric acid (1900). The total oxygen content was determined by the method of Oxygen (1900). The total hydrogen content was determined by the method of Hydrogen (1900). The total carbon content was determined by the method of Carbon (1900). The total nitrogen content was determined by the method of Nitrogen (1900). The total phosphorus content was determined by the method of Phosphorus (1900). The total sulfur content was determined by the method of Sulfur (1900). The total calcium content was determined by the method of Calcium (1900). The total magnesium content was determined by the method of Magnesium (1900). The total potassium content was determined by the method of Potassium (1900). The total sodium content was determined by the method of Sodium (1900). The total iron content was determined by the method of Iron (1900). The total copper content was determined by the method of Copper (1900). The total zinc content was determined by the method of Zinc (1900). The total manganese content was determined by the method of Manganese (1900). The total cobalt content was determined by the method of Cobalt (1900). The total nickel content was determined by the method of Nickel (1900). The total chromium content was determined by the method of Chromium (1900). The total boron content was determined by the method of Boron (1900). The total molybdenum content was determined by the method of Molybdenum (1900). The total selenium content was determined by the method of Selenium (1900). The total tellurium content was determined by the method of Tellurium (1900). The total iodine content was determined by the method of Iodine (1900). The total bromine content was determined by the method of Bromine (1900). The total fluorine content was determined by the method of Fluorine (1900). The total chlorine content was determined by the method of Chlorine (1900). The total oxygen content was determined by the method of Oxygen (1900). The total hydrogen content was determined by the method of Hydrogen (1900). The total carbon content was determined by the method of Carbon (1900).

ATVARADO VASQUEZ NICOLAS

SECRET

Noni's original position

REF ID: A68911

STRT TOROSHA OPERATIV


CONFIDENTIAL

80	400.000.-	400.000.-
20	100.000.-	100.000.-
1	5.000.-	5.000.-

5,000.00

5,000.00

5,000.00



Central Intelligence Agency

1-986-1

Edson Alvarado V.

SECRET

100-443887-100

[illegible]

NOTAS:

51, según el Art. 209 de la Ley de Compañías, la sociedad habiendo actualizado adecuadamente la información de la compañía en la columna "Apellidos y Nombres completos" y los valores correspondientes a las acciones.

readquiridas en las columnas respectivas

(1) Para inversionistas extranjeros

(2) Para inversión extranjera directa

(3) Para uso de Superintendencia
