



BAYVIEW SOCIAL- THERAPY, P.O. 5A.

[illegible]

NAME: _____ **DATE:** _____ **PERIOD:** _____ **PAGE NO.:** _____

[illegible]

1000-0000/01/0000-0000\$10.00/0

ACTIVATION OF CYP1A1 BY CYP1B1 IN CYP1A1-DEFICIENT CELLS

doi:10.1371/journal.pone.0142800.g002

Abstract

Ignacio QUARE, Carlos QUAREZ, Fernando Tello, Luisito VILLALBA, Juan A.
 FRANCISCO DE GONZALEZ, Roberto DE JESUS, JOSEPH VILLALBA, Carlos
 RODRIGUEZ, LAURENCE, RAY J. CHAVEZ, TONYA, JAMES VILLALBA, TONYA

Handwritten signature

1111

Author: WILLIAM L. JONES
Editor: JAMES H. JONES
Editor: JAMES H. JONES
Editor: JAMES H. JONES

Figure 1. A schematic diagram of the experimental design. The subjects were divided into two groups: the control group and the experimental group. The control group received a standard training program, while the experimental group received a modified training program. The results of the training program were compared between the two groups.

Hydroxide is a strong base
Structure is OH^-
Symbol is OH^-
IDEAL is 14