



Abstract

100



WOLFF, J. H. 1993. *Wolff's Catalog of Fishes*. American Fisheries Society, Bethesda, Maryland.

© 2004 Blackwell Publishing Ltd *Journal of Internal Medicine* 255: 111–118

1

100

Abstract



100

1000

100

100

© 2004 Blackwell Publishing Ltd, *Journal of Internal Medicine* 255: 103–110

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 399–406

100

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenolic content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total sterol content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenolic content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total sterol content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).

Table 1

[illegible]

Abstract

Year	Number of publications	Percentage of total
1990	1	1.0
1991	1	1.0
1992	1	1.0
1993	1	1.0
1994	1	1.0
1995	1	1.0
1996	1	1.0
1997	1	1.0
1998	1	1.0
1999	1	1.0
2000	1	1.0
2001	1	1.0
2002	1	1.0
2003	1	1.0
2004	1	1.0
2005	1	1.0
2006	1	1.0
2007	1	1.0
2008	1	1.0
2009	1	1.0
2010	1	1.0
2011	1	1.0
2012	1	1.0
2013	1	1.0
2014	1	1.0
2015	1	1.0
2016	1	1.0
2017	1	1.0
2018	1	1.0
2019	1	1.0
2020	1	1.0
2021	1	1.0
2022	1	1.0
2023	1	1.0
2024	1	1.0
2025	1	1.0
2026	1	1.0
2027	1	1.0
2028	1	1.0
2029	1	1.0
2030	1	1.0
2031	1	1.0
2032	1	1.0
2033	1	1.0
2034	1	1.0
2035	1	1.0
2036	1	1.0
2037	1	1.0
2038	1	1.0
2039	1	1.0
2040	1	1.0
2041	1	1.0
2042	1	1.0
2043	1	1.0
2044	1	1.0
2045	1	1.0
2046	1	1.0
2047	1	1.0
2048	1	1.0
2049	1	1.0
2050	1	1.0
2051	1	1.0
2052	1	1.0
2053	1	1.0
2054	1	1.0
2055	1	1.0
2056	1	1.0
2057	1	1.0
2058	1	1.0
2059	1	1.0
2060	1	1.0
2061	1	1.0
2062	1	1.0
2063	1	1.0
2064	1	1.0
2065	1	1.0
2066	1	1.0
2067	1	1.0
2068	1	1.0
2069	1	1.0
2070	1	1.0
2071	1	1.0
2072	1	1.0
2073	1	1.0
2074	1	1.0
2075	1	1.0
2076	1	1.0
2077	1	1.0
2078	1	1.0
2079	1	1.0
2080	1	1.0
2081	1	1.0
2082	1	1.0
2083	1	1.0
2084	1	1.0
2085	1	1.0
2086	1	1.0
2087	1	1.0
2088	1	1.0
2089	1	1.0
2090	1	1.0
2091	1	1.0
2092	1	1.0
2093	1	1.0
2094	1	1.0
2095	1	1.0
2096	1	1.0
2097	1	1.0
2098	1	1.0
2099	1	1.0
2100	1	1.0

RESEARCH DESIGN



100

... ..

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

Nome: _____

41

SRI@gob.ec

